Linear plus fractional multiobjective programming problem with homogeneous constraints using fuzzy approach

Sanjay Jain¹ and Kailash Lachhwani²

We develop an algorithm for the solution of multiobjective linear plus fractional programming problem (MOL+FPP) when some of the constraints are homogeneous in nature. Using homogeneous constraints, first we construct a transformation matrix T which transforms the given problem into another MOL+FPP with fewer constraints. Then, a relationship between these two problems, ensuring that the solution of the original problem can be recovered from the solution of the transformed problem, is established. We repeat this process of transformation until all the homogeneous constraints are removed. Then, we discuss the multi objective programming part, for which fuzzy programming methodology is proposed which works for the minimization of perpendicular distances between two hyper planes (curves) at the optimal points of the supremum perpendicular distance. A compromised optimal solution is obtained as a result of the minimization of the The supremum perpendicular distance. The supremum perpendicular distance is obtained using the transformation matrix. Finally, an example is given to illustrate the proposed model.

Key words: Linear plus fractional functional, Transformation matrix, Distance function, *Membership function*.

1. Introduction

Decision-making problems occuring in the real world are generally multi criteria decision-making (MCDM) problems. Many researchers such as Wallenius [12], Hanan [6], Feng [4], Chanas [1] and Rommelfanger [10] used and or modified the concept of decision making in fuzzy environment. They discussed different approaches to deal with the multiobjective programming problems. Also, many iterative algorithms have been studied as given by Charnes and Cooper [2], Swarup [11], Martos [9], Jain and Mangal [7] for solving fractional programming problems. Jain and Lachhwani [8] discussed a solution methodology for multiobjective linear fractional programming problem, in

which the objective function in of the form $\frac{f(X)}{g(X)}$. Here, we address form which we denote by

MOL+FPP.

A multiobjective linear plus fractional programming problem (MOL+FPP) seeks to optimize more than one objective functions of the form $f(X) + \frac{g(X)}{h(X)}^{-1}$, that is the sum of a linear and a

quotient function, where f(X), g(X) and h(X) are linear functions subject to linear

¹Department of Mathematics, Government College, Ajmer, India, E-mail: <u>drjainsanjay@gmail.com</u>

² Government Engineering College, Bikaner, India, E-mail: <u>kailashclachhwani@yahoo.com</u>

(1)

(2)

constraints under the assumptions that the set of the feasible solutions is a convex polyhedron with a finite number of extreme points and that the denominators of the fractional part of the objective functions are nonzero on the feasible set.

We present an algorithm that performs to be better than other iterative methods as it needs computing time for the optimization process and obtains an optimum solution for a set of objective functions of such a linear plus fractional form.

The remainder of the paper is organized as follows. In Section 2, a transformation matrix T is constructed. In the Section 3, the relationship between the original problem and the transformed problem is given. In Section 4, we develop fuzzy programming model for the problem by minimizing the perpendicular distances between two hyper planes $Z_i(x) = \overline{Z_i}$, and $Z_i(x) = \overline{Z_i}$ where $\overline{Z_i}$ and $\overline{Z_i}$ are the maximum and minimum values of the function $Z_i(x)$ in the feasible region, respectively. Suitable membership functions are defined and a compromise optimal solution is obtained. In Section 5, a pseudo program is given for the construction of the transformation matrix T. An example is worked out in Section 6 to explain the model. Particular cases and conclusions are given in Section s 7 ad 8, respectively.

Maximize $\{Z_1(X), Z_2(X), ..., Z_k(X)\}$

2. Development of transformation matrix

Consider the problem,

subject and

where,

$$\begin{aligned} &Z_i(X) = \left(L_i X + l_{0i}\right) + \frac{C_i X + c_{oi}}{D_i X + d_{oi}} & \forall i = 1, 2, ..., k \end{aligned}$$
 to
$$\begin{aligned} &AX = b, \\ &X \ge 0 \end{aligned}$$

with $a_{i1}x_1 + a_{i2}x_2 + \dots + a_{in}x_n = 0$, for some *i*.

Here, L_i , C_i and D_i (i = 1, 2, ..., k) are row vectors with *n* components, *X* and *b* are column vectors with *n* and *m* components, respectively, $A = (A_1, A_2, ..., A_n)$ is an *m* by *n* matrix and the c_{oi} , d_{oi} (i = 1, 2, ..., k) are scalars. It is assumed that $D_i X + d_{oi} > 0$, over L, where $L = \{X : AX = b, X \ge 0\}$. Let $X = (x_1, x_2, ..., x_n)$ be an efficient solution of (2). If x_k and $a_{ik} > 0$, then it is obvious that there exists at least one x_l with $a_{il} < 0$. Taking this in view, we partition matrix A as $A = \{A^0, A^+, A^-\}$, where A^0 is the set of all columns of A, whenever $a_{ij} = 0$, (Let the number of such columns be r), A^+ be the set of all columns of A, whenever $a_{ij} > 0$, p in number, and finally A^- be the set of all columns of A, whenever $a_{ij} < 0$, q in number. Thus, p + q + r = n.

Now, we define a transformation matrix T (n by pq + r) such that the i-th equation of ATw = b will be automatically zero. Here, w is a column vector with pq + r components. This is accomplished by defining w_{kl} for each pair (k, l) such that $A_k \in A^+$ and $A_l \in A^-$.

Now, we partition $T = (T_1, T_2)$ where T_1 consists of unit column vectors e_j corresponding to w_{kl} . A column corresponding to the w_{kl} looks like:

$$egin{aligned} 0 \ -a_{ij} \ 0 \ a_{ik} \ 0 \end{bmatrix}.$$

Thus, T, which has *n* rows and r + pq columns, can be represented as:

$$T = (T_1, T_2) = [\langle e_j \rangle \forall j \in a_{ij} = 0; \langle t_{kl} \rangle \forall k \in A^+ \land l \in A^-],$$

that is, e_j is the *j* th column of the identity matrix I_n , and $t_{kl} = -a_{il}e_k + a_{ik}e_l$. (3) A pseudo code program is also framed for the construction of transformation matrix T in Section 5.

Theorem 2.1 In matrix T of order $n \times (pq + r)$, the i – th equation of ATw = b will be identically zero.

Proof: To prove this theorem, it is sufficient to show that the *i*-th equation of AT will have all the zero. Let A^i denote the *i*-th row of A. For any $j \in A^0$, it is clear that $A^i e_j = 0$, implying thereby that $A^i T_1 = 0$. Again for any (k, l), $k \in A^+$, and $l \in A^-$, we have $A^i t_{kl} = A^i (-a_{il}e_k + a_{ik}e_l) = -a_{il}a_{ik} + a_{ik}a_{il} = 0$. Thus, the left hand side of the *i*-th equation of AT will be zero and $b_i = 0$.

3. Transformed problem

Using the transformation X = Tw, we define the following problem corresponding to problem (2):

Maximize
$$z_i = \left(\overline{L_i} \mathbf{w} + l_{0i}\right) + \frac{C_i \mathbf{w} + c_{oi}}{\overline{D_i} \mathbf{w} + d_{oi}} \quad \forall i = 1, 2, ..., k$$
 (4)
Subject to $\overline{A} \mathbf{w} = b$
and $\mathbf{w} \ge 0$. (5)

In view of Theorem 2.1, the *i*-th equation in system (5) will be identically zero and hence can be removed while solving the program (4)-(5). The following theorems are now in order.

Theorem 3.1 If X solves (2), then there exists w(X=Tw) which solves (5).

Proof: Here, constraints and non-negativity restrictions are similar in both original and the transformed problem for the single objective to generalized problem taken by Chadha [3]. So, the proof can be seen in Chadha [3].

Theorem 3.2 If X^* solves the program (1)-(2), then $w^*(X^*=Tw^*)$ solves the program (4)-(5).

Proof: Theorem 3.1 guarantees the existence of a feasible w^* , i.e., $\overline{A}w^* = b$, $w^* \ge 0$. Next, since X* solves the program (1)-(2), then

$$\left(L_{i}X^{*}+l_{0i}\right)+\frac{C_{i}X^{*}+c_{0i}}{D_{i}X^{*}+d_{0i}} \geq \left(L_{i}X+l_{0i}\right)+\frac{C_{i}X+c_{0i}}{D_{i}X+d_{0i}}, \ \forall \ X \in L, \ i=1,2,...,k$$

Or equivalently,

$$\left(L_{i} \mathrm{Tw}^{*} + l_{0i}\right) + \frac{C_{i} \mathrm{Tw}^{*} + c_{0i}}{D_{i} \mathrm{Tw}^{*} + d_{0i}} \ge \left(L_{i} \mathrm{Tw} + l_{0i}\right) + \frac{C_{i} \mathrm{Tw} + c_{0i}}{D_{i} \mathrm{Tw} + d_{0i}}, \ \forall \ \mathbf{w} \in G, \ i = 1, 2, ..., k ;$$

or,

$$\left(\overline{L_i}\mathbf{w}^* + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w}^* + c_{0i}}{\overline{D_i}\mathbf{w}^* + d_{0i}} \ge \left(\overline{L_i}\mathbf{w} + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w} + c_{0i}}{\overline{D_i}\mathbf{w} + d_{0i}}, \ \forall \ \mathbf{w} \in G;$$

Thus, w^* solves the program (4)-(5).

Theorem 3.3 If w^* solves the program (4)-(5), then there exists $X^*=Tw^*$ which solves the program (1)-(2), and the extreme values of the two objective functions are equal. **Proof:** w^* being a solution of the program (4)-(5) means that

$$\overline{A}w^* = b, w \ge 0;$$

*

or

$$ATw^{*} = b, \text{ or } AX^{*} = b, \tag{6}$$

Furthermore, $T \ge 0$, $w^* \ge 0$, imply that $X^* \ge 0$ (7) Also, we know that,

*

$$\left(\overline{L_i}\mathbf{w}^* + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w}^* + c_{0i}}{\overline{D_i}\mathbf{w}^* + d_{0i}} \ge \left(\overline{L_i}\mathbf{w} + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w} + c_{0i}}{\overline{D_i}\mathbf{w} + d_{0i}}, \quad \forall \mathbf{w} \in G, \ i = 1, 2, \dots, k$$
(8)

And,

$$\left(L_{i}X^{*}+l_{0i}\right)+\frac{C_{i}X^{*}+c_{0i}}{D_{i}X^{*}+d_{0i}} \ge \left(L_{i}X+l_{0i}\right)+\frac{C_{i}X+c_{0i}}{D_{i}X+d_{0i}}, \ \forall \ X \in L, \ i=1,2,...,k$$
(9)

For a contradictory argument, let \overline{X} and not X^{*} solve the program (4)-(5), which means that

$$\left(L_{i}\overline{X}+l_{0i}\right)+\frac{C_{i}X+c_{0i}}{D_{i}\overline{X}+d_{0i}} \ge \left(L_{i}X^{*}+l_{0i}\right)+\frac{C_{i}X^{*}+c_{0i}}{D_{i}X^{*}+d_{0i}}, \qquad i=1,2,...,k$$

From Theorem 3.1 it follows that

$$\left(L_{i}T\overline{w} + l_{0i}\right) + \frac{C_{i}Tw + c_{0i}}{D_{i}T\overline{w} + d_{0i}} > \left(L_{i}Tw^{*} + l_{0i}\right) + \frac{C_{i}Tw^{*} + c_{0i}}{D_{i}Tw^{*} + d_{0i}}, \text{ for any } i$$

or,

$$\left(\overline{L_i}\mathbf{w} + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w} + c_{0i}}{\overline{D_i}\mathbf{w} + d_{0i}} \ge \left(\overline{L_i}\mathbf{w}^* + l_{0i}\right) + \frac{\overline{C_i}\mathbf{w}^* + c_{0i}}{\overline{D_i}\mathbf{w}^* + d_{0i}}.$$

This violates (8) and the contradiction proves the result. Finally, let Z_i^* and z_i^* be the optimal values of the objective functions of (1) and (4) at X* and w*, respectively. This means

$$Z_{i}^{*} = (L_{i}X^{*} + l_{0i}) + \frac{C_{i}X^{*} + c_{0i}}{D_{i}X^{*} + d_{0i}}$$
$$= (L_{i}Tw^{*} + l_{0i}) + \frac{C_{i}Tw^{*} + c_{0i}}{D_{i}Tw^{*} + d_{0i}}$$
$$= (\overline{L_{i}}w^{*} + l_{0i}) + \frac{\overline{C_{i}}w^{*} + c_{0i}}{\overline{D_{i}}w^{*} + d_{0i}} = z_{i}^{*} \quad \forall \ i = 1, 2, ..., k.$$
(10)

The result then follows from (5), (6), (8) and (9).

4. Solution of linear plus fractional multiobjective programming problem

Using the transformation X = Tw, the problem (1)-(2) reduces to:

Maximize $\{z_1(w), z_2(w), ..., z_k(w)\}$

Linear plus fractional multiobjective programming

where

and

where,
$$z_i(\mathbf{w}) = (L_i \mathbf{w} + l_{0i}) + \frac{\partial I}{\overline{D}_i \mathbf{w} + d_{oi}}$$
 $\forall i = 1, 2, ..., k$
subject to $\overline{A}\mathbf{w} = b$
and $\mathbf{w} \ge 0$. (11)

 $(\overline{-}, \overline{-}, \overline{-$

Problem (11) does not contain homogeneous constraints. This MOFLPP can be reduced to a fuzzy programming problem. This model is developed to minimize the perpendicular distances between two parallel hyperplanes $z_i(w) = \overline{z_i}$ and $z_i(w) = \overline{z_i}$, where $\overline{z_i}$ and z_i are the maximum and minimum values of the objective function $z_i(w)$. We define the distance function d with unit weight as:

$$d_i(\mathbf{w}) = \left|\overline{z_i} - z_i(\mathbf{w})\right| \qquad \forall i = 1, 2, ..., k$$

This distance depends upon w. At w = w (ideal point in w-space) as, given by Gupta and Chakraborty [5], d = 0 and at $w = \underline{w}$ (nadir point in w-space), $z_i(w) = z_i$, and we get the maximum value of $d_i(w)$ as:

$$\overline{d_i} = \left| \overline{z_i} - \underline{z_i} \right| \quad \forall \ i = 1, 2, \dots, k .$$
(12)

Treating this criterion to be of equal importance, the vector maximum problem (11) may be modeled as follows:

Find an action $w \in G$, which minimizes

$$\operatorname{Max}\left\{\left|\overline{z_{i}}-z_{i}(\mathbf{w})\right|, \ i=1,2,...,k\right\},\$$

$$G=\left\{\mathbf{w}:\overline{A}\mathbf{w}=b, \ \mathbf{w}\geq0\right\},$$
(13)

where,

We define the membership $\mu_i(d_i(\mathbf{w}))$ as follows:

$$\mu_{i}\left(d_{i}\left(\mathbf{w}\right)\right) = \begin{cases} 0 & \text{if } \mathbf{d}_{i}\left(\mathbf{w}\right) \geq p \\ \frac{p - d_{i}\left(\mathbf{w}\right)}{p} & \text{if } 0 < d_{i}\left(\mathbf{w}\right) < p \\ 1 & \text{if } \mathbf{d}_{i}\left(\mathbf{w}\right) \leq 0 \end{cases}$$

where, $p = \sup \left\{ \overline{d_i} \right\} \quad \forall \quad i = 1, 2, 3, ..., k$.

If λ be the minimum of all $\mu_i(d_i(\mathbf{w}))$, then,

$$d_i(\mathbf{w}) \leq -p\lambda + p$$
, *i.e.* $\overline{z_i} - z_i(\mathbf{w}) \leq -p\lambda + p$.

Now, the problem reduces to:

Maxλ

subject to

and

$$-z_{i}(\mathbf{w}) + p.\lambda \leq p - \overline{z_{i}}, \qquad \forall \ i = 1, 2, ..., k$$
$$\overline{A}\mathbf{w} = b$$
$$\lambda, \ \mathbf{w} \geq 0$$
(14)

which can be solved by non-linear programming techniques.

5. Pseudo code program for transformation matrix

The matrix T can be obtained using a program in C, C++ or any other programming language. For construction of this program, a pseudo code program of the procedure is as follows.

 $/\!/$ m is the number. of rows of matrix, n is the number. of columns.

// A [1: m] [1: n] is the two dimensional matrix with m rows and n columns.

// P is the number of possible elements of matrix A in the i-th row.

// Q is the number of negative elements of matrix A in the i-th row.

// R is the total number. of columns in the transformation matrix.

 $// e_j$ is the j-the column of the identity matrix

// e_j is the j-the colum
{
FOR I: = 1 to m do
FOR J: = 1 to n do
READ A [1: m][1:n];
READ I;
// FOR Ax=b where b=0:
FOR J: = 1 to n do
{
IF (A [I][J]=0) then
{
T[R] = e_j;
R=R+1;
}
FOR J: =1 to n do
{
IF (A [I] [J] < 0) then
}
</pre>

P: =P+1; X [P]: = J;

IF (A [I] [J] >0) then { Q: =Q+1; Y [Q] = J;

}
}
FOR K: =1 to P do
{
FOR L: =1 to Q do
{
T[R] =- a [I] [Y [L]]* e_[K]+ a [I] [Y [K]]* e_{Y[L]};
R: = R+1:
}
WRITE T [1: R]; // output transformation matrix.

6. Numerical example

Example. The following example explains the algorithm.

Maximize $\{Z_1(X), Z_2(X)\}$ where, $Z_1(X) = 2x_1 + \frac{2x_1 + 6x_2}{x_1 + x_2 + 1}$, and $Z_2(X) = 2x_2 + \frac{x_1 + x_2}{x_1 - x_2 + 1}$ subject to, $x_1 + x_2 + x_3 = 4$ $3x_1 + x_2 - x_4 = 6$ Linear plus fractional multiobjective programming

and

$$\begin{aligned} x_1 - x_2 + 0.x_3 + 0.x_4 &= 0\\ x_1, x_2, x_3, x_4 \ge 0 \,. \end{aligned}$$

The solution procedure using our proposed methodology can be well explained stepwise, as given in tabular form Table 1.

 Table 1. Solution of example 1

	The equivalent problem without homogeneous constraints is given by
Step 1.	Maximize $\{z_1(w), z_2(w)\}$
	$(-)$ \overline{C} W CTW $8W$
	where $z_1 = (L_1 w + l_{01}) + \frac{C_1 w}{\overline{D_1} w + 1} = (LTw + l_{01}) + \frac{CTw}{DTw + 1} = 2w_3 + \frac{3w_3}{2w_3 + 1}$
	and $z_2 = 4w_3$
	subject to $\overline{A}w = b$ or $ATw = b$,
	and $w \ge 0$
	Using the given pseudo code and consequently constructed C-program, we obtain:
Step 2.	
	$T = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}$
	0 1 0
	Using transformation matrix T, our problem becomes:
Step 3.	Maximize $\begin{cases} 2w + \frac{8w_3}{4w} \end{cases}$
	$2w_3 + 2w_3 + 1, 1w_3$
	subject to $w_1 + 2w_3 = 4$
	$w_2 + 4w_3 = 6$
	and $w_1, w_2, w_3 \ge 0$
	over region $G = \{ w : \overline{A}w = b, w \ge 0 \}$.
Step 4.	Here, we find $\overline{z_1} = 6$ $\underline{z_1} = 0$, $\overline{z_2} = 6$ $\underline{z_2} = 0$ and $p = \sup\{\overline{d_i}\} = 6$.
	Then, the given problem reduces into vector maximum problem,
Step 5.	Max. λ
	subject to $6\lambda \le 2w_2 + \frac{8w_3}{2}$
	$(2w_3 + 1)$
	$6\lambda \le 4w_3$
	$w_1 + 2w_3 = 4$
	$-w_2 + 4w_3 = 6$
	and $w_1, w_2, w_3, \lambda \ge 0$
	Solving this nonlinear programming problem, the compromise optimal solution of
Step 6.	transformed problem is obtained as:

$$w_{1} = 0, w_{2} = 2, w_{3} = 2, \lambda = \frac{16}{15}.$$

The solution of the original problem (X* = Tw) is given as:
$$\begin{bmatrix} x_{1} \\ x_{2} \\ x_{4} \\ x_{5} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 0 \\ 2 \end{bmatrix},$$

That is, the compromise optimal solution of (2.1) is:
$$x_{1} = 2, x_{2} = 2, x_{3} = 0, x_{4} = 2 \text{ and } Z_{1}^{*} = \frac{36}{5}, Z_{2}^{*} = 8.$$

7. Particular cases

1. If we take $L_i = 0$, $\forall i = 1, 2, ..., k$, and take only a single objective function with homogeneous constraints, then our problem is reduced to LFPP. This discussion is also given by Chadha [3]. 2. If we take $D_i = 0$, $\forall i = 1, 2, ..., k$, then $\overline{D_i} = 0$, and it our problem is reduced to a MOLPP. This discussion is also given by Gupta and Chakraborty [5], by defining the distance function

 $d_i(\mathbf{w})$ with the weight $\frac{1}{\left\{\sum c_{ij}^2\right\}^{1/2}}$



Figure 1. Selection of ideal point

8. Conclusions

When an MOFPP has more than one homogeneous constraints, then we_can extend our approach for its solution. In general, T(s) is determined only when AT(1), AT(2),...,AT(s-1), have been computed. This algorithm reduces the number of constraints.

Linear plus fractional multiobjective programming

Compromised solution depends on the choice of nadir point (lowest justifiable value) of the objective function. When the justifiable value changes, the compromised solution also changes. As seen in Figure, if Z is the ideal point and N and N' we the two different minimum aspiration levels then their compromised solution are P and P', respectively, because NZ and N'Z are the direction in which the decision parameter λ is increased. In our methodology, to find minimum aspiration level, we used minimum value of each objective function. This point is the ideal point for the vector minimization problem of the same objective functions with the same constraints, which generally lies outside the feasible region. Knowing the nadir point (worst point), and zenith point (ideal point) we can find the direction of the decision parameter λ in which λ is increased. Considering the region by taking the lowest justifiable value each objective function gets equal importance in the optimization process.

Reference

- [1] Chanas, S. (1989), Fuzzy programming in multi objective linear programming a parametric approach, *Fuzzy sets and system*, 29, 303 313.
 - [2] Charnes, A. and Cooper, W.W., (1962), Programming with linear fractional unctional, *Naval Research log.Quart.*,9, 181-186.
 - [3] Chadha, S.S. (1999), A linear fractional program with homogeneous constraints, *OPSEARCH*, 36, 390-398.
 - [4] Feng, Y.J. (1983), A method using fuzzy mathematical programming to solve vector maximum problem, *Fuzzy sets and systems*, 9, 129 136.
 - [5] Gupta, S. and Chakraborty, M. (1997), Multi objective linear programming: A fuzzy programming approach, *International journal of management and system*, 13(2), 207 214.
 - [6] Hanan, E.L. (1981), Linear programming with multiple goals, *Fuzzy sets and systems*, 6, 235-248.
 - [7] Jain, S. and Mangal, A. (2006), Solution of a multi objective fractional programming problem, *Journal of Indian Academy of Mathematics*, 28(1), 133 141.
 - [8] Jain, S. and Lachhwani, K. (2009) "Multi objective fractional programming problem: Fuzzy programming approach", *Proceedings of National Academy of Sciences, India-Physical sciences (Section- A)*, Vol. 79, part III.
 - [9] Martos, B. (1975), *Non-linear Programming, Theory and methods*, North-Holland Publishing Company, Amsterdam.
 - [10] Rommelfanger, H. (1989), Interactive decision making in fuzzy Linear Optimization problems, *European Journal of Operational Research*, 41, 210 217.
 - [11] Swarup, K. (1965), Linear Fractional Functional Programming, *Operations Research*, 13, 1029-1036.
 - [12] Wallenius, J. (1975), Comparative evaluation of some interactive approaches to multicriteria optimization, *Management science*, 21, 1387-1396.

Downloaded from iors.ir on 2025-08-26