ABS SOLUTION OF NORMAL EQUATIONS OF SECOND KIND AND APPLICATION TO THE PRIMAL-DUAL INTERIOR POINT METHOD FOR LINEAR PROGRAMMING

Emilio Spedicato, Marco Bonomi and Antonino Del Popolo Department of Mathematics, University of Bergamo, Bergamo, Italy Email: emilio@unibg.it

Abstract

We consider on application of the ABS procedure to the linear systems arising from the primal-dual interior point method where Newton methods is used to compute a path to the solution. When approaching the solution the linear system, which has the form of normal equations of the second kind, becomes more and more ill conditioned. We show how the use of the Huang algorithm in the ABS class can reduce the ill conditioning. Preliminary numerical experiments show that the proposed approach can provide a residual in the computed solution up to sixteen orders lower.

Key words : ABS methods, normal equations of the second kind, Huang algorithm, primal-dual interior point method, Newton method.

1. Introduction

ABS algorithms have been introduced in 1984 by Abaffy, Broyden and Spedicato [1] to solve determined or underdetermined linear systems and have been later extended to linear least squares, nonlinear equations, optimization problems and integer (Diophantine) equations and LP problems. The ABS literature consists presently of over 400 papers; see Abaffy and Spedicato's monograph [2] for a presentation of such techniques as available at the beginning of 1989 and Spedicato et al. [6,7] for reviews dealing with later results. The class of ABS methods unifies most existing methods for solving linear systems and provides a variety of alternative ways of implementing a specific algorithm. Extensive computational experience has shown that ABS methods are implementable in a stable way, being often more accurate than the corresponding traditional algorithms, and that on vector/parallel machines they often perform faster; see for example Bodon [3,4].

The linear programming problem is one of the most important problems in optimization, both in the continuous and discrete forms. The classical algorithm is the simplex method, developed by Dantzig following a suggestion by Von Neumann. This

method approaches the solution by moving over the vertices of the polytope defining the feasible region. The simplex method, while usually performing efficiently, may in the worst case require an exponentially high cost, since all vertices may have to be reached. About a quarter of a century ago the new class of interior point methods has been developed where the iterates were generated along a path lying inside the boundary of the polytope, with possible computational advantages. These algorithms have been widely investigated in close to five thousand papers. The most efficient implementation appears to be the one based upon the primal-dual approach, leading to solving a sequence of structured nonlinear equations, containing partly linear and partly nonlinear equations in a diagonal form. The linear system to be solved at each iteration of Newton's method has the form of normal equations of the second kind, that is,

$AA^{T}x = b$

where $x \in R^n, b \in R^m, m < n$, and the matrix A has the form

$$A = BD^{-1}$$

where B is m by n and D is the diagonal matrix whose i - th element in approching the solution tends either to zero or to infinity, resulting in a very ill conditioned matrix A and an even more ill conditioned matrix AA^{T} .

The normal equations of the second kind arise when one is looking for the unique solution of least Euclidean norm of an underdetermined system. In the context of ABS methods, such equations can be solved in a way similar to that proposed for the normal equations of the first kind, (see [2]), without forming the product AA^{T} , thereby reducing the condition number essentially to the square root. Moreover, the diagonal matrix responsible for the ill conditioning can be removed from the matrix being applied to the right hand side. This will be discussed in the next section. The last section presents preliminary results of the proposed approach, which show that extremely better accuracy in the residual error can be obtained than by working on the system in the original form.

For reader's nvenience, we recall the basic steps of the (unscaled) ABS class for solving the linear system Ax = b, $x \in R^n$, $b \in R^m$, $m \le n$, $A = (a_1, \dots, a_m)^T$.

(A) Set $x_1 \in \mathbb{R}^n$ arbitrary, $H_1 \in \mathbb{R}^{n,n}$ arbitrary nonsingular, i = 1.

Algorithm 1: The unsculed ABS algorithm.

(B) Compute the vector $s_1 = H_i a_i$ and the scalar $\tau_i = a_i^T x_i - b_i$. If $s_i \neq 0$ then go to (C), the i –th equation is linearly independent from the previous equations. If $\tau_i \neq 0$ then stop, the system has no solution, else remove $\tau_i = 0$ the i –th equation, being a linear combination of the previous ones, and set $x_{i+1} = x_i$, $H_{i+1} = H_i$ and go to (F). (C) Compute the search direction $p_i = H_i^T z_i$, $z_i \in \mathbb{R}^n$ arbitrary save that $z_i^T H_i a_i \neq 0$.

(D) Update the estimate of the solution by $x_{i+1} = x_i - \alpha_i p_i$, with $\alpha_i = (a_i^T x_i - b_i)/a_i^T p_i$.

(E) Update the (Abaffian) matrix H_i by $H_{i+1} = H_i - H_i a_i w_i^T H_i / w_i^T H_i a_i$, where $w_i \in \mathbb{R}^n$ is arbitrary save that $w_i^T H_i a_i \neq 0$.

(F) If i = m, then stop x_{m+1} solves the system, else increment i by one and go to (B).

In the above class (that can be generalized to the scaled ABS class by introducing an extra parameter, obtaining a complete realization of any Krylov- type process computing the solution in no more than m steps), algorithms are defined by specific choices of the parameters I, z_i and w_i . For the present problem, we use the so called Huang or implicit Gram-Schmidt algorithm, defined by the choices

$$H_1 = I, z_i = w_i = a_i$$
,

where a_i i the transpose 8 the i – th row of A. This algorithm has the remarkable property that it determines the unique solution of least Euclidean norm of an underdetermined system, if it is started with an arbitrary vector x_1 proportional to a_1 , usually the zero vector. The solution is moreover approached, if $x_1 = 0$, monotonically from below. Additionally, the search directions p_i are orthogonal among themselves. The algorithm's stability can be improved in several ways. Usually one does a reprojection on the search direction, that is.

$$\mathbf{p}_{i} = \mathbf{H}_{i}(\mathbf{H}_{i}\mathbf{a}_{i})$$

and defines the update, Un an example as

$$\mathbf{H}_{i+1} = \mathbf{H}_i - \mathbf{p}_i \mathbf{p}_i^{\mathrm{T}} / \mathbf{p}_i^{\mathrm{T}} \mathbf{p}_i \; .$$

The above modification of the Huang algorithm is called the modified Huang algorithm. For the use of the ABS algorithm in the normal equations of the second kind step (B) plays an important role, since it allows the elimination of the redundant equations and in practice often of the equations that are almost linearly dependent. Notice that in practice the norms of s_i and τ_i are not tested against zero, due to the effects of roundoff, but against some positive small ϵ . The best determination of ϵ is still an open problem, but in practice we have found in all our extensive testing of ABS algorithm that ϵ can be taken rather safely as the product of the machine zero by the norm of the tested quantity.

2. ABS solution of the normal equations of the second kind

It is immediate to see that the normal equations of the second kind are equivalent to the following extended system, where only A appears,

 $A^{\mathrm{T}}x = y$ Ay = b.

The first of the above equations is an overdetermined linear system that must be

solvable. Hence y must lie in the column space of the coefficient matrix A^{T} , i.e, in the row space of the matrix of the second system, which is an underdetermined system. Now, an underdetermined system has one and only one solution in the row space, which is the solution of smallest Euclidean norm. Such a solution is computed by the Huang algorithm started with the zero vector. Therefore, we are led to the following procedure, apparently never considered before, for solving the given normal equations of the second kind.

Algorithms: Solution for normal equations 8th Second kind.

AA - Solve by Huang algorithm, started with the zero vector, system Ay = b.

BB - Solve by any ABS algorithm the overdetermined system $A^{T}x = y$, where the n - m linearly dependent equations are identified and removed at step (B) of algorithm 1.

By Algorithm 2 we deal with systems having only matrix A, not the conditionally "squared" matrix AA^{T} , reducing the condition number. We also notice that if A is not of full rank, then the Huang algorithm will still compute the solution of Ay = b of smallest Euclidean norm, while use of the Huang algorithm in the overdetermined system, which is in such a case compatible but with infinite solutions, would compute the solution of smallest Euclidean norm.

With such a solution, Newton method would be well defined and convergent under mild conditions. Not many years ago, exactly using this property of Huang algorithm we eliminated occasional nonconvergence of Newton method used to solve the equations arising from the simulation of a large plant system defined by algebraic and differential equations. Our contractor, ENEL, sold the stabilized code to its French counterpart for a substantial amount.

Now we pay more attention to the case arising from the primal-dual interior point method, where A has the form $A = BD^{-1}$. The extended system is now written as

 $D^{-1}B^Tx \,=\, y$

$$BD^{-1}y = b.$$

In the primal-dual method the matrix B is fixed, D changes at every iteration. Since and D^{-1} scales the columns of the underdetermined system, the solution will require different search vectors at every Newton iteration. Once y is computed, we notice that the overdetermined system can be written in the form

 $B^{T}x = Dy$.

Since the coefficient matrix B is unchanged in the course of Newton's iterations, the following can be done only once at the first iteration:

AAA - Computation of the search vectors.

BBB - Identification of the redundant equations, which are exactly those equations associated with dependent rows in B^{T} , since A and B have exactly the same set of dependent/independent equations. Therefore, only the diagonal components of D associated with the selected independent rows of B are used in the computation of the solution (all of them enter in the underdetermined system).

Hence, it is enough to determine independent equations and the associated search vectors (which do not depend on D) only at the first Newton is iteration. This approach, in other term is equivalent to essentially dealing with the linear part of the primal-dual equations only once at the beginning. The number of operations to solve the overdetermined system after the first step is of $O(m^2)$. The number of operations to solve the underdetermined system by Huang algorithm is of $O(mn^2)$. These estimates can be certainly reduced when A is sparse, albeit a full study of the best implementation of a sparse Huang algorithm (possibly via the vector-wise formulation that applies to all ABS algorithms) is yet to be presented.

3. Some numerical experiments

Here we present a few numerical experiments obtained on two types of matrices:

1 - The ill conditioned Micchelli matrix where $A_{i,j}$ is the difference i - j in absolute value. Such a matrix is symmetric nonsingular with the condition number growing fast with the dimension. It appears in radial basis function approximation.

2 - A well conditioned Householder matrix with truncated last row and postmultiplied by a diagonal matrix with elements tending alternatively to zero and infinity. This was done through multiplication by the diagonal matrix whose elements had the form alternatively 10^k , 10^{-k} , for k = 1, 2, 3, 4, 5. The values taken for n were n = 10, 100, 200, for m were m = 2, n/4, n/2, 3n/4, n - 2. The right hand side of the original equation was computed by setting all components of x equal to one. Notice that this nominal solution is the one computed by the algorithm in exact arithmetic only for full rank problems; for rank deficient problems, or problems with deficient numerical ranks as determined in step (B) of Algorithm 1, the Huang algorithm would determine the solution of smallest Euclidean norm, generally different from the nominal solution. The proposed algorithm via the extended system and use of the modified Huang algorithm is compared in the following experiments with the results of applying the modified Huang algorithm directly on the original system, $AA^Tx = b$. Now we give some results obtained in double precision, where E1, E2 are the errors in the solution respectively by using the original system and the extended system and R1, R2 are similarly the residual errors, both computed in the Euclidean norm. For more results, see Bonomi [5].

m = 2	n = 200	E1 = 4.7E - 12	E2 = 6.8E - 11	R1 = 0	R2 = 5E - 13
m = n/4	n = 200	E1 = 2.3E - 7	E2 = 7.9E - 7	R1 = 1.7E - 7	R2 = 4E-11
m = n/2	n = 200	E1 = 5.7E - 7	E2 = 1.5E - 6	R1 = 4.8E - 7	R2 =6.2E-11
m = 3n/4	n = 200	E1 = 8.9E - 7	E1.6 = E - 7	R1 = 7.7E - 7	R2 =1.4E-10
m = n - 2	n = 200	E1 = 1.9E - 6	E2 = 2.1E - 6	R1 = 1.3E - 6	R2 =1E-10

Table 1. Results with the Micchelli matrix

Table 2. Results with the Householder matrix, scaled by diagonal matrix with $\epsilon = 10^{-1}$

m = 2	n = 200	E1 = 0	E2 = 1E - 15	$\mathbf{R}1=0$	R2 = 3E - 15
m = n/4	n = 200	E1 = 2.3E - 13	E2 = 5.7E - 13	R1 = 2.3E - 13	R2 =2E-14
m = n/2	n = 200	E1 = 4.2E - 12	E2 = 50E - 12	R1 = 4.8E - 13	R2 =5.3E-14
m = 3n/4	n = 200	E1 = 8.9E - 12	E1.6E – 11	R1 = 3.7E - 13	R2 = 4.2E-14
m = n - 2	n = 200	E1 = 4.7E - 11	E2 = 5.4E - 11	R1 = 9.5E - 13	R2 =1E-13

Table 3. Results with the Householder matrix, scaled by diagonal matrix with $\epsilon = 10^{-5}$

m = 2	n = 200	E1 = 2E - 16	E2 = 2E - 15	R1 = 1.8E - 12	R2 = 1.4E - 11
m = n/4	n = 200	E1 = 2.4	rankdef	R1 = 2.1E - 5	R2 =8E-9
m = n/2	n = 200	E1 = 6.3	rankdef	R1 = 4.7E + 10	R2 =2E-6
m = 3n/4	n = 200	E1 = 9	rankdef	R1 = 5.1E + 10	R2 =6.7E-4
m = n - 2	n = 200	E1 = 10	rankdef	R1 = 3.8E + 10	R2 =6E-5

Inspecting the above Tables, we propose the following preliminary conclusions: A - On the Micchelli matrix, we see little difference in the solution error, while the

algorithm using the extended system is up to four orders better in the residual error. B - There is a clear difference in performance between the two algorithms when we consider the ill conditioned matrix obtained by scaling the Householder matrix. The algorithm based on the extended system is up to 16 orders better in the residual error, while no comparison can be made in the solution error since the matrices turn out to be numerically rank deficient, so that the nominal solution is not generally the least Euclidean norm solution. There is little difference when the matrix, scaled by $\epsilon = 1/10$, is still well conditioned.

C - The accuracy obtained by solving the extended system tends to increase with m approaching n.

Further experiments are planned on several types of matrices, providing also estimates of the condition number and of the running time, that are not given in this case.

Acknowledgements Work done in the framework of a research supported by Bergamo University Academic Funds 2007. This work originated during a visit in Bergamo by prof. Florian Potra of the University of Maryland, supported by GNCS.

Refrence

bibitem 1. Abaffy, J., Broyden C. G. and Spedicato, E., A class of direct methods for linear systems, Numer. Math. 45, 361-376, 1984.

2. Abaffy, J. and Spedicato, E., ABS Projection Algorithms: Mathematical Techniques for Linear and Nonlinear Equations, Ellis Horwood, Chichester, 1989.

3. Spedicato, E., Bodon E., DEl Popolo A. and Xia Z., ABS algorithms for linear systems and optimization: a review and a bibliography, Ricerca Operativa 89, 39- 88, 2000

4. Bodon E, Numerical experiments with ABS algorithms on banded systems of linear equations, Report TR/PA/93/13, CERFACS, Toulouse, 1993

5. Bodon E, Numerical experiments with ABS algorithms on upper-banded systems of linear equations, Report TR/PA/93/14, CERFACS, Toulouse, 1993

6. Bonomi M., Utilizzo dei metodi ABS nei metodi interior point, Tesi di Laurea, AA 2006/07, Bergamo University. 2007

bibitem 7. Spedicato, E., Bodon, E., A. Del Popolo and Mahdavi-Amiri, N., ABS methods and ABSPACK for linear Systems and optimization: A review, 40R L, 51-66.

This document was created with Win2PDF available at http://www.daneprairie.com. The unregistered version of Win2PDF is for evaluation or non-commercial use only.