

On the Behavior of Damped Quasi-Newton Methods for Unconstrained Optimization

M. Al-Baali^{1,*}, L. Grandinetti²

We consider a family of damped quasi-Newton methods for solving unconstrained optimization problems. This family resembles that of Broyden with line searches, except that the change in gradients is replaced by a certain hybrid vector before updating the current Hessian approximation. This damped technique modifies the Hessian approximations so that they are maintained sufficiently positive definite. Hence, the objective function is reduced sufficiently on each iteration. The recent result that the damped technique maintains the global and superlinear convergence properties of a restricted class of quasi-Newton methods for convex functions is tested on a set of standard unconstrained optimization problems. The behavior of the methods is studied on the basis of the numerical results required to solve these test problems. It is shown that the damped technique improves the performance of quasi-Newton methods substantially in some robust cases (as the BFGS method) and significantly in certain inefficient cases (as the DFP method).

Keywords: *Unconstrained optimization, Quasi-Newton methods, Line-search framework.*

Manuscript received on 22/10/2011 and accepted for publication on 31/12/2011.†

1. Introduction

We study the behavior of the recent class of damped quasi-Newton methods, proposed by Al-Baali [7] for solving the unconstrained optimization problem

$$\min_{x \in R} f(x),$$

where f is a nonlinear differentiable function. This damped (D-) class resembles that of Broyden with line searches (see, for example, Dennis and Schnabel [11], Fletcher [12] or Nocedal and Wright [26]) except that the change in gradients $\gamma_k = g_{k+1} - g_k$ is replaced by the hybrid damped-technique

$$\hat{\gamma}_k = \varphi_k \gamma_k + (1 - \varphi_k) B_k \delta_k, \quad (1)$$

where $\varphi_k \in (0, 1]$ is a parameter, before updating a Hessian approximation B_k . Here, $g_k = \nabla f(x_k)$, $B_k \approx \nabla^2 f(x_k)$, $\delta_k = x_{k+1} - x_k$ and x_k is the current estimate of a solution of the problem.

†Invited paper.

*Corresponding Author.

¹ Department of Mathematics and Statistics, Sultan Qaboos University, Muscat, Oman. Email: albaali@squ.edu.om.

² Department of Electronics, Informatics and Systems, Calabria University, Rende 87036, Italy. Email: lugran@unical.it.

We notice that the value of $\varphi_k = 1$ (or $\widehat{\gamma}_k = \gamma_k$) reduces the damped class of methods to the Broyden family of methods. The latter family contains the standard BFGS and DFP methods, while the former class contains for some $\varphi_k \neq 1$ the corresponding D-BFGS and D-DFP methods, respectively. The D-BFGS method was applied to unconstrained optimization problems for the first time by Al-Baali [4, 5] who extended the D-BFGS method of Powell [21] for constrained optimization in augmented Lagrange and SQP methods (for further detail on the latter case, see for example Fletcher [12] Nocedal and Wright [20] or Gill and Leonard [14]).

Although the BFGS method is robust and has several useful numerical and theoretical properties, it suffers from a certain type of ill-conditioned problems. Therefore, several modification techniques have been introduced to the BFGS method see for example Yuan, [25] Zhang *et al.* [26] Li and Fukushima [16] Xu and Zhang [23] Zhang and Xu [27] Gill and Leonard [14] Al-Baali, [4,5] Wei *et al.* [22] Yabe *et al.* [24] Li *et al.* [17] Al-Baali and Khalfan [8] Al-Baali and Grandinetti [7] and the references therein). Since the latter paper also shows that the above D-technique is preferable to the other modifications for γ_k in the BFGS method, here we consider testing not only the D-technique when introduced not only to the inefficient DFP method, but also to other members of the Broyden family of methods. The remainder of our work is organized as follows. In Section 2, we describe the class of damped methods and consider some safeguarded schemes which maintain the useful theoretical and numerical properties of the BFGS method. Section 3 describes some numerical results obtained by applying a selection of methods to a set of standard test problems. It is shown that the proposed damped technique improves the performance of quasi-Newton method substantially in some robust cases (like the BFGS method) and significantly in certain inefficient cases (like the DFP method). Finally, Section 4 gives our concluding remarks. Sections 5 and 6 are appendices.

2. Damped Quasi-Newton Methods

Here we describe the D-Broyden class of quasi-Newton methods. At the beginning of each iteration, a positive definite Hessian approximation B_k is used to define the search direction s_k by solving the system of linear equations $B_k s = -g_k$. Then, a step-length α_k is chosen such that the following Wolfe-Powell conditions hold:

$$f_k - f_{k+1} \geq \sigma_0 \delta_k^T g_k \quad (2)$$

and

$$\delta_k^T \gamma_k \geq -(1 - \sigma_1) \delta_k^T g_k, \quad (3)$$

where $\sigma_0 \in (0, 0.5)$ and $\sigma_1 \in (\sigma_0, 1)$. Note that the latter inequality ensures that the curvature condition $\delta_k^T \gamma_k > 0$ holds so that the positive definiteness property holds for both the damped and ‘undamped’ Broyden class of methods. For the next iteration, B_k is updated to a new Hessian approximation,

$$B_{k+1} = B_k - \frac{B_k \delta_k \delta_k^T B_k}{\delta_k^T B_k \delta_k} + \frac{\widehat{\gamma}_k \widehat{\gamma}_k^T}{\delta_k^T \widehat{\gamma}_k} + \theta_k \widehat{w}_k \widehat{w}_k^T, \quad (4)$$

$$\widehat{w}_k = \left(\delta_k^T B_k \delta_k \right)^{1/2} \left(\frac{\widehat{\gamma}_k}{\gamma_k^T \delta_k} - \frac{B_k \delta_k}{\delta_k^T B_k \delta_k} \right), \quad (5)$$

where $\widehat{\gamma}_k$ is given by (1), and θ_k and φ_k are parameters. This class of damped methods is reduced to the well-known Broyden family of methods, if $\varphi_k = 1$ (or $\widehat{\gamma}_k = \gamma_k$), for all k , and to the BFGS and DFP methods, if, in addition, $\theta_k = 0$ and $\theta_k = 1$, respectively. The corresponding damped methods are referred to as D-BFGS and D-DFP, respectively.

The above D-Broyden methods is proposed by Al-Baali [7] by extending the D-BFGS method of Al-Baali [4, 5] for unconstrained optimization on the basis of the D-BFGS method of Powell [21] for constrained optimization. Al-Baali [5] uses the damped technique (1) with the choice

$$\varphi_k = \begin{cases} \frac{\sigma_2}{1 - \rho_k}, & \rho_k < 1 - \sigma_2, \\ \frac{\sigma_3}{\rho_k - 1}, & \rho_k > 1 + \sigma_3, \\ 1, & \text{otherwise,} \end{cases} \quad (6)$$

where

$$\rho_k = \frac{\gamma_k^T \delta_k}{\delta_k^T B_k \delta_k}, \quad (7)$$

$0 < \sigma_2 < 1$, and $\sigma_3 > 0$, which is reduced to that of Powell if $\sigma_2 = 0.8$ and $\sigma_3 = \infty$.

For sufficiently small values of σ_2 and σ_3 , the choice (6) ensures that $\widehat{\gamma}_k^T \delta_k$ is sufficiently close to the positive value of $\delta_k^T B_k \delta_k$ and, hence, the damped formula (4) maintains Hessian approximations sufficiently positive definite. However, the Broyden family satisfies this property only under the restrictions that $\gamma_k^T \delta_k > 0$ and $\theta_k > -1/a_k$, where

$$a_k = b_k h_k - 1, \quad b_k = \frac{\delta_k^T B_k \delta_k}{\delta_k^T \gamma_k}, \quad h_k = \frac{\gamma_k^T B_k^{-1} \gamma_k}{\delta_k^T \gamma_k}. \quad (8)$$

Since $a_k \geq 0$ (by the Cauchy inequality), the above useful property holds in particular for the nonnegative members $\theta_k = 0$ and $\theta_k = 1$. Another well-known member of the Broyden family is the symmetric rank 1 (SR1) update, defined by $\theta_k = 1/(1 - b_k)$, which does not belong to the convex class of updates. Since this update does not guarantee the above positive definiteness property and negative values of θ_k seem to work well in practice [28] Al-Baali [2] suggested the switching BFGS/SR1 update, given by the non-positive choice

$$\theta_k = \begin{cases} \frac{1}{1 - b_k}, & h_k < 1, \\ 0, & \text{otherwise.} \end{cases} \quad (9)$$

Although the corresponding method of (9) converges globally for convex objective functions, its performance is better than that of the robust BFGS method (see for instance Lukšan and Spedicato [18] and the next section). The latter two choices for θ_k also define the damped D-SR1 and D-(BFGS/SR1) updates, respectively. These damped updates satisfy the positive definiteness property for sufficiently small values of φ_k .

We now outline the damped-Broyden family of quasi-Newton methods.

Algorithm 2.1. Damped-Broyden Family

0. Give a starting point x_1 , a symmetric positive-definite initial Hessian approximation B_1 , values of σ_0 and σ_1 , and set $k := 1$.
1. Terminate if a convergence test holds.
2. Compute the search direction $s_k = -B_k^{-1}g_k$.
3. Find a step-length α_k and a new point $x_{k+1} = x_k + \alpha_k s_k$ such that the following strong Wolfe-Powell conditions hold:

$$f_{k+1} \leq f_k + \sigma_0 \alpha_k g_k^T s_k, \quad |g_{k+1}^T s_k| \leq -\sigma_1 g_k^T s_k. \quad (10)$$
4. Compute δ_k, γ_k and ρ_k .
5. Choose values for θ_k and φ_k and compute $\hat{\gamma}_k$.
6. Update B_k by the D-Broyden formula (4).
7. Set $k := k + 1$ and go to Step 1.

This algorithm is reduced to the normal Broyden family of methods if the choice $\varphi_k = 1$ is used in Step 5 for all iterations (which is also obtained by substituting $\sigma_2 = 1$ and $\sigma_3 = \infty$ into (6)). This choice with, in particular, $\theta_k = 0$ yield the standard BFGS method, while $\varphi_k \neq 1$, for some k , yields a D-BFGS method. We use in Step 3 the strong Wolfe-Powell conditions, as commonly used in practice, which imply the Wolfe-Powell conditions (2)-(3). Note that Al-Baali [7] also extends the global and superlinear convergence result of Byrd et al. [9], that a restricted Broyden family of methods has for convex functions, to the class of damped methods.

3. Numerical Analysis

In this section, we test the performance of Algorithm 2.1 for some values of the updating parameter θ_k , which was implemented, as in Al-Baali and Grandinetti [7] in Fortran 77, using the Lahey software with double precision arithmetic. In Step 0 of the algorithm, we let the initial Hessian approximation $B_1 = I$, the identity matrix, and use the values of $\sigma_0 = 10^{-4}$ and $\sigma_1 = 0.9$ in (10). The run was stopped in Step 1 when either

$$\|g_k\|^2 \leq \epsilon \max(1, |f_k|),$$

where ϵ is the machine epsilon ($\approx 10^{-16}$), $f_{k+1} \geq f_k$, or the number of iterations reached

10^5 . In Step 3, we used the scheme (2. 6. 4) of Fletcher [12] for obtaining an acceptable step-length α_k for the strong Wolfe-Powell conditions (10). This scheme is based on some function interpolations and firstly tries Fletcher's initial estimate (2. 6. 8) for α_k , which is reduced to one in the limit. In Step 5, some values for θ_k were considered as below. The default value of $\varphi_k=1$ was usually used, but for the damped technique is considered we let φ_k be defined by formula (6) with several values of σ_2 and σ_3 chosen on the basis of some results reported in Al-Baali [7] Here, we report the results for the following choices which differ from those considered by Al-Baali and Grandinetti [7] for the D-BFGS method. We let

$$\sigma_2 = \begin{cases} 0.5, & \rho_k < 0.5 \text{ and } |\theta_k| a_k > 0.5, \\ \max(\min(0.5, \bar{\sigma}), \nu), & \rho_k < 0.5 \text{ and } |\theta_k| a_k > 0.5, \\ 1, & \text{otherwise,} \end{cases} \quad (11)$$

where $\nu = 10^{-7}$ and

$$\bar{\sigma} = \frac{0.5|1-\rho_k|}{\sqrt{|\theta_k| a_k}}. \quad (12)$$

We also let σ_3 be given by (11) and (12) with $<$, 0.5, 1, and $|\theta_k|$ replaced by $>$, e , ∞ , and $\max(|\theta_k|, 1)$, respectively. The small value of $\nu = 10^{-7}$ was used to avoid destroying the character of the damped technique. Indeed, this value was never used in our experiments, since we observed that the smallest value for σ_2 and σ_3 was 10^{-5} , which rarely occurred in practice. Thus, we employed the damped technique when the values of the scalars $|1-\rho_k|$ and $b_k h_k - 1$ became sufficiently away from zero, because Al-Baali [7] showed that these scalars tend to zero and $\varphi_k \rightarrow 1$, when the damped methods converge to the solution superlinearly for convex functions.

To define the parameter θ_k in Step 5, we tried several selections for θ_k . Here, we report the results for the three well known choices of $\theta_k = 0$, $\theta_k = 1$ and (9), which maintain the positive definite Hessian approximations. These choices yield the BFGS, DFP and BFGS/SR1 and their corresponding D-BFGS, D-DFP and D-(BFGS/SR1) methods, respectively. We applied these methods (as in Al-Baali and Grandinetti, [7]) to a set of 89 standard test problems, with their names, citations and dimensions (in the range [2,100]) listed in Table 3 in Appendix B.

As expected, the DFP method was inefficient, since it failed to solve about 36% of the test problems and converged very slowly for several other test problems. However, the other methods solved all the test problems successfully.

To examine the behavior of the successful methods, the numerical results are summarized in tables 1 and 2. Table 1 represents the ratios of the total number of line searches, function evaluations and gradient evaluations required by each method to solve all the test problems in the set to that required by the BFGS method (denoted by T_l , T_f and T_g ,

respectively). These ratios clearly show that the damped methods are preferable to the undamped ones. They indicate that the total number of l , f and g evaluations required to solve all the tests in the set by the D-BFGS, D-(BFGS/SR1) and D-DFP methods are at most 57%, 62% and 76%, respectively, of those required by the BFGS method. Thus, the damped technique improves the performance of the BFGS method substantially and DFP method significantly.

Since the ratios in Table 1 do not adequately illustrate the performance of the methods, we also present Table 2. The column headings A_l , A_f and A_g stand for certain ‘average’ ratios related, respectively, to the number of l , f and g evaluations required to solve each test problems by the methods versus those required by the BFGS method, using the fair rule of Al-Baali (see for example Al-Baali [3] and Appendix B). A value of $A_l < 1$ (similarly for A_f and A_g) indicates that the performance of a method compared to that of BFGS is improved by $100(1 - A_l)\%$ in terms of the number of l .

Although the corresponding ratios for each method in Table 2 are larger than those in Table 1, these ratios maintain the following observations. The damped technique plays an important role for improving the performance of robust and inefficient quasi-Newton methods. We observe that the performance of the D-DFP method is a little better than the standard BFGS method, the other three methods perform substantially better than BFGS and D-BFGS is the most efficient method. The latter method performs about 24%, 17% and 23% better than the BFGS method in terms of the number of l , f and g evaluations, respectively. Although BFGS/SR1 performs much better than the BFGS method, D-(BFGS/SR1) also performs a little better than BFGS/SR1 in terms of l and g and slightly in terms of f . This observation indicates that the damped technique does not destroy the features of robust methods. We also note that the most efficient D-BFGS method is slightly better than the D-(BFGS/SR1) method.

Table 1. Ratios of total cost as compared to BFGS

| Method | T_l | T_f | T_g |
|--------------|-------|-------|-------|
| D-BFGS | 0.532 | 0.573 | 0.538 |
| D-DFP | 0.736 | 0.764 | 0.774 |
| BFGS/SR1 | 0.810 | 0.866 | 0.932 |
| D-(BFGS/SR1) | 0.552 | 0.615 | 0.579 |

Table 2. Average ratios as compared to BFGS

| Method | A_l | A_f | A_g |
|--------------|-------|-------|-------|
| D-BFGS | 0.763 | 0.826 | 0.767 |
| D-DFP | 0.924 | 0.971 | 0.936 |
| BFGS/SR1 | 0.841 | 0.888 | 0.872 |
| D-(BFGS/SR1) | 0.780 | 0.865 | 0.802 |

4. Conclusion

We showed that the damped technique works well in practice. The technique improves the performance of inefficient methods significantly and robust methods substantially. The D-BFGS method was recommended, although further experiments are required of finding typical values for σ_2 and σ_3 or other useful choices for the damped parameter φ_k . It is also worth introducing the self-scaling technique to the efficient damped methods in a manner similar to that of Al-Baali and Khalfan [8] who showed that combining the damped and self-scaling techniques yielded a substantial improvement of the BFGS method.

References

- [1] Al-Baali, M. (1991), A rule for comparing two methods in practical optimization, Research Report No. 119, Systems Department, Calabria University, Italy.
- [2] Al-Baali, M. (1993), Variational quasi-Newton methods for unconstrained optimization. *Journal of Optimization Theory and Applications*, 77, 127-143.
- [3] Al-Baali, M. (1994), An efficient class of switching type algorithms in the Broyden family, *Optimization Methods and Software*, 4, 29-46.
- [4] Al-Baali, M. (2003), Quasi-Wolfe conditions for Newton-like methods, Presented at 18th International Symposium on Mathematical Programming, Copenhagen, Denmark.
- [5] Al-Baali, M. (2004), Quasi-Wolfe conditions for quasi-Newton methods for large-scale optimization, Presented at 40th Workshop on Large Scale Nonlinear Optimization, Erice, Italy.
- [6] Al-Baali, M. (2009), Convergence properties of damped Broyden's family of quasi-Newton methods, Research Report DOMAS 09/1, Sultan Qaboos University, Oman.
- [7] Al-Baali, M. and Grandinetti, L. (2009), On practical modification of quasi-Newton BFGS method, *Advanced Modeling and Optimization*, 11, 63-76.
- [8] Al-Baali, M. and Khalfan, H. (2008), A combined class of self-scaling and modified quasi-Newton methods, Research Report DOMAS 08/1, Sultan Qaboos University, Oman (to appear in *Computational Optimization and Applications*).
- [9] Byrd, R.H., Liu, D.C. and Nocedal, J. (1992), on the behavior of Broyden's class of quasi-Newton methods, *Society for Industrial and Applied Mathematics Journal on Optimization*, 2, 533-557.
- [10] Conn, A.R., Gould, N.I.M. and Toint, Ph.L. (1988), Testing a class of algorithms for solving minimization problems with simple bound on the variables, *Mathematics of Computation*, 50, 399-430.
- [11] Dennis, J.E. and Schnabel, R.B. (1996), Numerical Methods for Unconstrained Optimization and Nonlinear Equations, Society for Industrial and Applied Mathematics, Philadelphia, USA.
- [12] Fletcher, R. (1987), Practical Methods of Optimization, Second Edition, Wiley, Chichester, England (reprinted in 2000).
- [13] Fletcher, R. and Powell, M.J.D. (1963), A rapidly convergent descent method for minimization, *Computer Journal*, 6, 163-168.

- [14] Gill, P.E. and Leonard, M.W. (2003), Limited-memory reduced-Hessian methods for large-scale unconstrained optimization, *Society for Industrial and Applied Mathematics Journal on Optimization*, 14, 380-401.
- [15] Grandinetti, L. (1984), Some investigations in a new algorithm for nonlinear optimization based on conic models of the objective function, *Journal of Optimization Theory and Applications*, 43, 1-21.
- [16] Li, D. and Fukushima, M. (2001), A modified BFGS method and its global convergence in nonconvex minimization, *Journal of Computational and Applied Mathematics*, 129, 15-35.
- [17] Li, D., Qi, L. and Roshchina, V. (2008), A new class of quasi-Newton updating formulas, *Optimization Methods and Software*, 23, 237-249.
- [18] Lukšan, L. and Spedicato, E. (2000), Variable metric methods for unconstrained optimization and nonlinear least squares, *Journal of Computational and Applied Mathematics*, 124, 61-95.
- [19] Moré, J.J., Garbow, B.S. and Hillstom, K.E. (1981), Testing unconstrained optimization software, *ACM Transactions on Mathematical Software*, 7, 17-41.
- [20] Nocedal, J. and Wright, S.J. (1999), Numerical Optimization, Springer, London.
- [21] Powell, M.J.D. (1978), Algorithms for nonlinear constraints that use Lagrange functions, *Mathematical Programming*, 14, 224-248.
- [22] Wei, Z., Yu, G., Yuan, G. and Lian, Z. (2004), The superlinear convergence of a modified BFGS-type method for unconstrained optimization, *Computational Optimization and Applications*, 29, 315-332.
- [23] Xu, C. and Zhang, J. (2001), a survey of quasi-Newton equations and quasi-Newton methods for optimization, *Annals of Operations Research*, 103, 213-234.
- [24] Yabe, H., Ogasawara, H. and Yoshino, M. (2007), Local and superlinear convergence of quasi-Newton methods based on modified secant conditions, *Journal of Computational and Applied Mathematics*, 205, 617-632.
- [25] Yuan, Y. (1991), A modified BFGS algorithm for unconstrained optimization, *Institute of Mathematics and Applications Journal of Numerical Analysis*, 11, 325-332.
- [26] Zhang, J.Z., Deng, N.Y. and Chen, L.H. (1999), Quasi-Newton equation and related methods for unconstrained optimization, *Journal of Optimization Theory and Applications*, 102, 147-167.
- [27] Zhang, J.Z. and Xu, C.X. (2001), Properties and numerical performance of quasi-Newton methods with modified quasi-Newton equations, *Journal of Computational and Applied Mathematics*, 137, 269-278.
- [28] Zhang, Y. and Tewarson, R.P. (1988), Quasi-Newton algorithms with updates from the preconvex part of Broyden's family, *Institute of Mathematics and Applications Journal of Numerical Analysis*, 8, 487-509.

Appendix A

We now describe the rule of Al-Baali [1], and also for example [3], for comparing two methods (say M1 and M2) on the basis a set of pair numbers, say p_i and q_i , for $i=1,2,\dots,m$, related to M1 and M2, respectively. In this paper, $m (=89)$ denotes the number of test problems and both p_i and q_i denote, for all i , either the number of line searches, function evaluations or gradient evaluations required to solve a test i by the M1 and M2 methods, respectively (the latter method is referred to BFGS and the former one to another method under comparison).

Al-Baali modifies the well-known average ratio $\frac{1}{m} \sum_{i=1}^m \frac{p_i}{q_i}$ to the following modified ‘average’ measure,

$$A_R = \frac{1}{m} \sum_{i=1}^m r_i,$$

where r_i has the form

$$r_i = \begin{cases} \frac{p_i}{q_i}, & p_i \leq q_i, \\ 2 - \frac{p_i}{q_i}, & \text{otherwise.} \end{cases}$$

It is assumed that $r_i = 1$ if both $p_i, q_i \rightarrow \infty$, which is also used in the following cases. If both M1 and M2 methods either failed or converged to two different solutions, for some test i , then we set $r_i = 1$ (i.e., $p_i = q_i$). Thus, the A_R ratio takes all kinds of terminations into account and always belongs to the interval $[0, 2]$. A value of $A_R \leq 1$ indicates that the M2 method reduces the cost of (i.e., improves over) M1 by $100(1 - A_R)\%$ (or equivalently it is $1/A_R$ times better than M1). If $A_R > 1$, then M1 is better as in the latter sense but with A_R replaced by $(2 - A_R)$. Note that if the inequality $p_i \leq q_i$ holds for all i , then A_R is reduced to the usual average of the m ratios (p_i / q_i) .

Appendix B

Here we present Table 3 consisting of some details on the set of test problems used in this paper. The first column consists of codes and numbers of the tests given in the original sources. One of these tests is proposed by Fletcher and Powell [13] another can be seen in Grandinetti [15] and the other tests have been collected and described by Moré, et al. [19] and Conn, et al. [10]. The second column of the table records the number of variables n et al. used for each function. We note that the dimensions of 59 test problems range from 2 to 30 and those of the remaining 30 test problems are either 40 or 100. The symbol \dagger indicates that the same test function is used again, but with the initial point multiplied by 100. The third column of the table consists of the function names.

Table 3. The set of test problems

| Test Code* | Dimension n | Function's name |
|------------|---------------------|--|
| MGH3 | 2 | Powell badly scaled |
| MGH4 | 2 | Brown badly scaled |
| MGH5 | 2 | Beale |
| MGH7 | 3† | Helical valley |
| MGH9 | 3 | Gaussian |
| MGH11 | 3 | Gulf research and development |
| MGH12 | 3 | Box three-dimensional |
| MGH14 | 4† | Wood |
| MGH16 | 4† | Brown and Dennis |
| MGH18 | 6 | Biggs Exp 6 |
| MGH20 | 6,9,12,20 | Watson |
| MGH21 | 2†,10†,20†, 40, 100 | Extended Rosenbrock |
| MGH22 | 4†,12†,20†, 40, 100 | Extended Powell singular |
| MGH23 | 10,20, 40, 100 | Penalty I |
| MGH25 | 10†,20†,40, 100 | Variably dimensioned |
| MGH26 | 10,20, 40, 100 | Trigonometric of Spedicato |
| MGH35 | 8,9,10,20, 40, 100 | Chebysquad |
| CGT1 | 8 | Generalized Rosenbrock |
| CGT2 | 25 | Another chained Rosenbrock |
| CGT4 | 20 | Generalized Powell singular |
| CGT5 | 20 | Another generalized Powell singular |
| CGT10 | 30, 40, 100 | Toint's seven-diagonal generalization of Broyden tridiagonal |
| CGT11 | 30, 40, 100 | Generalized Broyden tridiagonal |
| CGT12 | 30, 40, 100 | Generalized Broyden banded |
| CGT13 | 30, 40, 100 | Another generalized Broyden banded |
| CGT14 | 30, 40, 100 | Another Toint's seven-diagonal of Broyden tridiagonal |
| CGT15 | 10 | Nazareth |
| CGT16 | 30, 40, 100 | Trigonometric |
| CGT17 | 8, 40, 100 | Generalized Cragg and Levy |
| CH-ROS | 10†,20†,40, 100 | Chained Rosenbrock |
| TRIGFP | 10,20, 40, 100 | Trigonometric of Fletcher and Powell |

* MGH m : Collected by Moré et al. [19] where m denotes the number of the problem test

CGT m : Collected by Conn et al. [10] where m denotes the number of the problem test

CH-ROS: Given by Grandinetti [15]

TRIGFP: Given by Fletcher and Powell [13].

†: Two initial points were used; the standard point \bar{x} and $100\bar{x}$.